



HEMOLIFE (Diagnostic Division of HEMOPET)

11561 Salinaz Avenue, Garden Grove, CA 92843

Tel: 714-891-2022; Fax: 714-891-2123

www.hemopet.org www.nutriscan.org E-Mail: info@hemopet.org

Status: FINAL REPORT
Notes: food intolerance, scratching, soft stool
Diet: Raw

Accession No. Test146503	Doctor SAMPLE REPORT	Owner SAMPLE REPORT	Pet Name SAMPLE REPORT	Received 06/25/20
Species Canine	Breed Golden Retriever	Sex FS	Pet Age 5 Yrs 55 lbs	Reported 07/08/20

Test Requested	Result	Case Specific	General Range	Units
Beef Salivary IgA	6.550 Negative Reaction		< 10	U/mL
Beef Salivary IgM	10.650 Weak Reaction		< 10	U/mL
Chicken Salivary IgA	8.745 Negative Reaction		< 10	U/mL
Chicken Salivary IgM	9.640 Negative Reaction		< 10	U/mL
Corn Salivary IgA	10.732 Weak Reaction		< 10	U/mL
Corn Salivary IgM	11.666 Borderline Reaction; Avoid		< 10	U/mL
Duck Salivary IgA	9.188 Negative Reaction		< 10	U/mL
Duck Salivary IgM	10.521 Weak Reaction		< 10	U/mL
Lamb Salivary IgA	8.480 Negative Reaction		< 10	U/mL
Lamb Salivary IgM	9.024 Negative Reaction		< 10	U/mL
Milk Salivary IgA	10.868 Weak Reaction		< 10	U/mL
Milk Salivary IgM	10.284 Weak Reaction		< 10	U/mL
Pork Salivary IgA	9.500 Negative Reaction		< 10	U/mL
Pork Salivary IgM	12.600 Intermediate reaction, Avoid		< 10	U/mL
Soy Salivary IgA	9.439 Negative Reaction		< 10	U/mL
Soy Salivary IgM	9.933 Negative Reaction		< 10	U/mL
Turkey Salivary IgA	10.184 Weak Reaction		< 10	U/mL
Turkey Salivary IgM	10.942 Weak Reaction		< 10	U/mL
Venison Salivary IgA	10.189 Weak Reaction		< 10	U/mL
Venison Salivary IgM	10.443 Weak Reaction		< 10	U/mL
Wheat Salivary IgA	10.199 Weak Reaction		< 10	U/mL
Wheat Salivary IgM	10.090 Weak Reaction		< 10	U/mL
White Fish Salivary IgA	11.140 Weak Reaction		< 10	U/mL
White Fish Salivary IgM	11.220 Weak Reaction		< 10	U/mL
Barley Salivary IgA	8.350 Negative Reaction		< 10	U/mL



ENRICHING PET LIFE THROUGH DIAGNOSTIC TESTING
world-wide personal mail-in diagnostic lab for vets and pets



HEMOLIFE (Diagnostic Division of HEMOPET)

11561 Salinaz Avenue, Garden Grove, CA 92843

Tel: 714-891-2022; Fax: 714-891-2123

www.hemopet.org www.nutriscan.org E-Mail: info@hemopet.org

Accession No. Test146503	Doctor SAMPLE REPORT	Owner SAMPLE REPORT	Pet Name SAMPLE REPORT	Received 06/25/20
------------------------------------	-------------------------	-----------------------------------	---------------------------	----------------------

Test Requested	Result	Case Specific	General Range	Units
Barley Salivary IgM	11.900 Borderline Reaction; Avoid		< 10	U/mL
Egg Salivary IgA	7.490 Negative Reaction		< 10	U/mL
Egg Salivary IgM	7.840 Negative Reaction		< 10	U/mL
Lentil Salivary IgA	10.104 Weak Reaction		< 10	U/mL
Lentil Salivary IgM	11.875 Borderline Reaction; Avoid		< 10	U/mL
Millet Salivary IgA	9.978 Negative Reaction		< 10	U/mL
Millet Salivary IgM	10.334 Weak Reaction		< 10	U/mL
Oatmeal Salivary IgA	10.900 Weak Reaction		< 10	U/mL
Oatmeal Salivary IgM	10.600 Weak Reaction		< 10	U/mL
Peanut Salivary IgA	8.699 Negative Reaction		< 10	U/mL
Peanut Salivary IgM	11.020 Weak Reaction		< 10	U/mL
Potato Salivary IgA	12.500 Intermediate reaction, Avoid		< 10	U/mL
Potato Salivary IgM	13.029 Medium Reaction; Avoid		< 10	U/mL
Quinoa Salivary IgA	10.294 Weak Reaction		< 10	U/mL
Quinoa Salivary IgM	10.416 Weak Reaction		< 10	U/mL
Rabbit Salivary IgA	8.590 Negative Reaction		< 10	U/mL
Rabbit Salivary IgM	9.435 Negative Reaction		< 10	U/mL
Rice Salivary IgA	11.000 Weak Reaction		< 10	U/mL
Rice Salivary IgM	10.750 Weak Reaction		< 10	U/mL
Salmon Salivary IgA	9.750 Negative Reaction		< 10	U/mL
Salmon Salivary IgM	9.375 Negative Reaction		< 10	U/mL
Sweet Potato Salivary IgA	10.413 Weak Reaction		< 10	U/mL
Sweet Potato Salivary IgM	9.266 Negative Reaction		< 10	U/mL



Dear Colleague and Owner: Nutriscan Panels #1 and 2 are completed and showed food reactions to: Corn (and Cornstarch in supplements), Pork, Barley (a gluten), Lentils (including Peas), and Potatoes. Please avoid all these reactive foods. Best wishes, Jean [W.Jean Dodds, DVM]

[Click here for a guide to read the Nutriscan results and additional foods derived from the 24 foods tested.](#)



HEMOLIFE (Diagnostic Division of HEMOPET)

11561 Salinaz Avenue, Garden Grove, CA 92843
Tel: 714-891-2022; Fax: 714-891-2123
www.hemopet.org www.nutriscan.org E-Mail: info@hemopet.org

Accession No. Test146503	Doctor SAMPLE REPORT	Owner SAMPLE REPORT	Pet Name SAMPLE REPORT	Received 06/25/20
------------------------------------	-------------------------	-----------------------------------	---------------------------	----------------------

Nutriscan test results indicate which of the 24 foods tested are unsuitable for this animal (i.e. tested or above 11.5 units/ml with *either* IgA or IgM antibodies).

Foods testing below 11.5 units/mL should be safe to feed as results were negative or weak (weak reactions reflect background assay "noise").

A **NutriGold** consultation is available, if you have questions or wish advice and suggestions about these results.

Note: Beef reactivity includes Buffalo and Bison; Chicken includes Chicken fat; Corn reactivity includes Cornstarch in supplements and treats; Cow Milk includes Cheese and Whey; Lamb reactivity includes Goat and Sheep and Goat dairy products; Pork includes bacon; White-Colored Fish reactivity includes sardines, herring and tuna (and their oils); Lentils includes Peas and Pea Protein.

Nutriscan Interpretation (For more information see [nutriscan](#))

Pet should not consume food or treats containing ingredient(s) showing results of **11.5 or greater, that state "Avoid" above, for either IgA and/or IgM antibodies.** Please remember that changing your pet's diet can cause bowel upsets, so gradually changing foods over 7-14 days is preferred. Recommend rechecking salivary food sensitivity or intolerance levels every 6 -12 months.

Degree of reactivity

- **<10 U/mL** indicates a normal food antigen tolerance level =**negative result.**
- **10-11.4 U/mL** indicates a weak reaction; **clinical significance unclear**
- **11.5-11.9 U/mL** indicates a **borderline reaction**
- **12-12.9 U/mL** indicates an **intermediate reaction**
- **13-14.9 U/mL** indicates a **medium reaction**
- **>= 15 U/mL** indicates a **strong reaction**

Differences between antibodies to IgA and IgM in Saliva

Antibodies to IgA measure the immune response to certain foods in secretions like saliva that have occurred over the last 2 years.



ENRICHING PET LIFE THROUGH DIAGNOSTIC TESTING
world-wide personal mail-in diagnostic lab for vets and pets



HEMOLIFE (Diagnostic Division of HEMOPET)

11561 Salinaz Avenue, Garden Grove, CA 92843
Tel: 714-891-2022; Fax: 714-891-2123
www.hemopet.org www.nutriscan.org E-Mail: info@hemopet.org

Accession No. Test146503	Doctor SAMPLE REPORT	Owner SAMPLE REPORT	Pet Name SAMPLE REPORT	Received 06/25/20
------------------------------------	-------------------------	-----------------------------------	---------------------------	----------------------

Antibodies to IgM measure the body's primary immune response to a recent exposure of certain foods within the last 6 months.

Recipes When Some Meats or Fish are Non-Reactive

Suggestions for the diet would be to include only non-reactive meat or fish; vegetables and fruit, with the possible addition of tapioca (cassava root = gluten-free starch). Suggest trying: Chick peas (garbanzo beans), pinto or kidney beans, carrots, zucchini, spinach, green beans, **but no peas or pea fiber (if Lentils are reactive)**, and blueberries, cranberries, pomegranates, (**avoid strawberries**); apples, pears, bananas, and melons (including watermelon). Then, if doing better, try adding tapioca. The dog may also tolerate quail, pheasant, kangaroo, or emu.

If Beef is non-reactive, Buffalo and Bison should be acceptable.

If Lamb is non-reactive, Goat should be OK (including Goat Milk & Yogurt).

Oils – When White-colored Fish and/or Salmon are Reactive

Sources of the important Omega-3 oils are: No menhaden, pollack, sardine or herring oil, if white-colored fish is reactive. No salmon oil, if salmon is reactive.

Krill oil = ancient crustacean (but not with a shellfish allergy).

Plant oils = olive oil, borage oil, sunflower oil, coconut oil (in moderation), primrose oil, moringa oil, but **not canola oil**.

Now that you have your Nutriscan results, consider doing CellBIO, our novel biomarker test to ascertain the risk for obesity, inflammation, infection, passive cigarette smoke exposure, and even cancer.

For more information see [cellbio](#).

Liz Jean Doores, DVM

HEMOLIFE assays and reports use patented non-RIA technology and interpretations. Thyroid 5™ is a trademark.